

Possible Solution to the Poincare Conjecture

Gordon Chalmers

e-mail: gordon@quartz.shango.com

Abstract

The Poincare conjecture is analyzed in the context of Calabi-Yau n -folds. A simple treatment is given by embedding the three-manifolds into these CY manifolds, and then taking the orbifold limit. The higher-dimensional proofs are also available in this context.

The Poincare conjecture states essentially that a simply connected three-manifold is homeomorphic to the S^3 . The proofs in four and higher dimensions are contained in [1] and [2]. The analagous statement for non-simply connected three manifolds is that they are homeomorphic to the S^3/Γ , with Γ a discrete subgroup of $SO(4)$.

A possible proof here is proffered by embedding the metric components of the 3-manifold into a Ricci-flat Calabi-Yau manifold. The Calabi-Yau manifolds have orbifold limits into T^{2n}/Z_p ; as such the metric components become functions of the orbifold in the smooth degeneration of the Calabi-Yau into the orbifold. The curvature of the metric solution on the tori are constant, hence models a sphere.

Consider the 6 metric components of a Riemannian 3-manifold, $g_{\mu\nu}$. Take these metric components as holomorphic coordinates of a Calabi-Yau 6-fold, $X = g$. The copy of the metric coordinates models the anti-holomorphic coordinates, $\bar{X}$. The set of points generated by the 3-manifold's metrics then span a locus of points in the Calabi-Yau 6-fold, possibly the entire holomorphic half of the 6-fold. The curvature of the metric on the space of g on a T^3/Γ has curvature zero, and the blow up is found by a diffeomorphism effectively which does not change the curvature condition. Several patches are required to avoid unphysical singularities, with zero curvature (one reason is by self-consistency as the space is homeomorphic to a T^3/Γ). The embedding is chosen by using the full metric (periodic) basis of the 3-manifold. In the case of manifolds with higher homotopy, the singularity of the 3 manifold is placed at the center of one of the CY 2-spheres in order to coincide with the orbifold limit of S^3/Γ , i.e. all the 2-spheres line up.

The curvature condition of the individual Calabi-Yau manifolds is $R = 0$. However, since the coordinates of these manifolds are the metric, the translation of this curvature into the 3-manifold's curvature is complicated,

$$R(y) = \Lambda(y) , \tag{1}$$

with $\Lambda(y)$ a non-negative function. This can be seen to be non-negative after the fact of taking the orbifold limit of the Calabi-Yau. A diffeomorphism of the 3-manifold is required to map the non-negative curvature into the form $\Lambda(y)$, together with the appropriate choice of the Calabi-Yau 6-fold.

The Calabi-Yau manifolds have orbifold limits of the form, $M_6 \rightarrow T^{12}/Z_p$, holomorphically. As the metric components g of the 3-manifold are the coordinates of the Calabi-Yau, in the smooth orbifold limit they span a locus of coordinates of the

T^{12}/Z_p . The toriodal limit produces the individual curvatures of the metric components as if they were on circles, and they follow as coordinates on $Z(y) = Z(y + 2\pi R)$.

The constant circle curvatures of these deformed metric components model that of a 3-manifold with constant curvature in all directions in the metric components. The question is how this is related to a 3-sphere. The holomorphic space parametrized by such metric components $X = g$ has the topology of a T^3/Z_q , with zero curvature, but with constant curvature which is non-vanishing along the circles. This has the same topology as a 3-sphere. Furthermore, the homotopy of the original 3-manifold is trivial, this requires the coordinates resulting from metric components, to be also trivial.

The original curvature condition of the 3-manifold follows from (1). A diffeomorphism of the 3 manifold can change this non-negative curvature into other functions $\tilde{\Lambda}(y)$. Or rather in the converse, what is required is to change the original 3-manifold's curvature into the form of (1). This curvature condition depends on the specific Calabi-Yau, in finding the conditions $R_6 = 0$ and $R_3(y) = \Lambda(y)$.

Thus holomorphically embedding the components of the 3-manifold into the 6-fold, together with the known smooth degenerations of the latter, generate the homeomorphisms of the former into the 3-spheres. The proof can also be run backwards, by starting with a 3-sphere and blowing it up into the Calabi-Yau, after including its anti-holomorphic half.

The same argument goes through for higher dimensional simply connected manifolds. Furthermore, the argument seems to show how Thurston's conjecture holds, for non-simply connected 3-manifolds and S^3/Γ .

Gordon Chalmers thanks Albert Whitehead of the Royal Theological Society of Newark.

References

- [1] M.H. Freedman, *The topology of four-dimensional manifolds*, J. Diff. Geom **17** (1982), 357-453.
- [2] S. Smale, *Generalized Poincare's conjecture in dimensions greater than four*, Annals. Math. **74** (1961) 391-406; Bull. Amer. Math. Soc. **66** (1960) 373-375.